

COMPOSITION AND PACKAGING:

Balm on the basis of medicinal plants in original bottles of 100 ml. With a protective seal, equipped with a measuring cap:

- 1) cardboard packaging with a protective branded holographic sticker;
- 2) abstract;

Lagohilus (*Lagochilus inebrians*); Nettles (*Urtica dioica* L.); Shepherd's bag (*Capsella bursa*); Water pepper (*Poligonum hydropiper* L.); Yarrow (*Achillea millefolium* L.); Taternik (*Onopordum acanthium*); Highland Serpent (*Poligonum bistorta*).

CHEMICAL COMPOSITION:

Phytoncides, flavonoids (quercetin, quercetrine, rutin), lagohilin, vitamins A, D, E, K, C, formic, pantothenic acid, carotenoids, essential oils, tannins.

CLINICAL EFFECTS:

- Haemostatic - increases the coagulation ability of blood due to the activation of plasma and cellular clotting factors;
- Antifibrinolytic - fibrinolysis proactivators are suppressed;
- Reduces the permeability of small vessels, small damaged vessels are clogged;
- Stimulates the contraction of the smooth muscles of the uterus;
- Stimulates erythropoiesis.

INDICATIONS FOR ADMINISTRATION:

- Uterine, gastrointestinal, pulmonary, hemorrhoidal hemorrhages;
- Diseases associated with increased permeability of blood vessels (hemorrhagic diathesis);
- Improvement of uterine contractility;
- To prevent increased bleeding during surgical interventions.

DOSAGE AND ADMINISTRATION:

For 5-10 ml. 3 times a day before meals, until the clinical effect is achieved. If necessary, you can dissolve in 50-100 ml. Tea, juice or in warm boiled water.

CONTRAINDICATIONS:

Increased coagulability of blood, thrombosis, thrombophlebitis, intestinal obstruction, hepatic coma. Individual intolerance of individual components of the drug.

SIDE EFFECTS:

There may be an allergic reaction to individual components.

DRUG INTERACTION:

In the appointment with other drugs, no negative effects were noted.

STORAGE CONDITIONS:

Store in a dry place, protected from light, out of reach of children, at a temperature of + 2 ° C to + 25 ° C;

During storage, cloudiness of the solution may be observed, which does not affect the activity of the preparation;

Shelf life is 24 months.